arXiv:nlin/0203031v1 [nlin.CD] 14 Mar 2002

The Lyapunov stability of first order dynamic equations with respect to time-dependent Riemannian metrics. An example.

Gennadi Sardanashvily[1]

Department of Theoretical Physics, Physics Faculty, Moscow State University, 117234 Moscow, Russia

Abstract. A simple example that I have been requested illustrates the statement in the E-print nlin.CD/0201060 that solutions of a smooth first order dynamic equation can be made Lyapunov stable at will by the choice of an appropriate time-dependent Riemannian metric.

Let $\mathbb{R}$ be the time axis provided with the Cartesian coordinate t. In geometric terms, a (smooth) first order dynamic equation in non-autonomous mechanics is defined as a vector field γ on a smooth fibre bundle $Y \to \mathbb{R}$ which obeys the condition $\gamma\rfloor dt = 1$. With respect to bundle coordinates (t, y^k) on Y, this vector field reads

$$\gamma = \partial_t + \gamma^k \partial_k. \tag{1}$$

The associated first order dynamic equation takes the form

$$\dot{y}^k = \gamma^k(t, y^j)\partial_k,$$

where $(t, y^k, \dot{t}, \dot{y}^k)$ are holonomic coordinates on the tangent bundle TY of Y. Its solutions are trajectories of the vector field γ (1).

Let a fibre bundle $Y \to \mathbb{R}$ be provided with a Riemannian fibre metric g, defined as a section of the symmetrized tensor product $\overset{2}{\vee} V^*Y \to Y$ of the vertical cotangent bundle V^*Y of $Y \to \mathbb{R}$. With respect to the holonomic coordinates $(t, y^k, \overline{y}_k)$ on V^*Y, it takes the coordinate form

$$g = \frac{1}{2} g_{ij}(t, y^k)\overline{d}y^i \vee \overline{d}y^j,$$

where $\{\overline{d}y^i\}$ are the holonomic fibre bases for V^*Y.

With a Riemannian fibre metric g, the instantwise distance $\rho_t(s, s')$ between two solutions s and s' of a dynamic equation γ on Y at an instant t is defined as the distance between the points $s(t)$ and $s'(t)$ with respect to the Riemannian metric $g(t)$ on the fibre

---

[1] E-mail address: sard@grav.phys.msu.su

Y_t over $t \in \mathbb{R}$. Then the (upper) Lyapunov exponent of a solution s' with respect to a solution s is the limit

$$K(s,s') = \overline{\lim_{t\to\infty}} \frac{1}{t} \ln(\rho_t(s,s')). \tag{2}$$

If the upper limit

$$\overline{\lim_{\rho_{t=0}(s,s')\to 0}} K(s,s') = \lambda$$

is negative, the solution s is said to be exponentially Lyapunov stable. If there exists at least one positive Lyapunov exponent, one speaks about chaos in a dynamical system.

Proposition 6 in E-print nlin.CD/0201060 states the following.

- Let λ be a real number. Given a dynamic equation defined by a complete vector field γ (1), there exists a Riemannian fibre metric on Y such that the Lyapunov spectrum of any solution of γ is λ.

The following example aims to illustrate this fact.

Let us consider one-dimensional motion on the axis $\mathbb{R}$ defined by the first order dynamic equation

$$\dot{y} = y \tag{3}$$

on the fibre bundle $Y = \mathbb{R}\times\mathbb{R} \to \mathbb{R}$ coordinated by (t, y). Solutions of the equation (3) read

$$s(t) = c\exp(t), \qquad c = \text{const.} \tag{4}$$

Let $e_{yy} = 1$ be the standard Euclidean metric on $\mathbb{R}$. With respect to this metric, the instantwise distance between two arbitrary solutions

$$s(t) = c\exp(t), \qquad s'(t) = c'\exp(t) \tag{5}$$

of the equation (3) is

$$\rho_t(s,s')_e = |c - c'|\exp(t).$$

Hence, the Lyapunov exponent $K(s,s')$ (2) equals 1, and so is the Lyapunov spectrum of any solution (4) of the first order dynamic equation (3).

Let now λ be an arbitrary real number. In accordance with Proposition 1 in E-print nlin.CD/0201060, there exists a coordinate

$$y' = y\exp(-t)$$

on $\mathbb{R}$ such that, written relative to this coordinate, the solutions (4) of the equation (3) read $s(t)$ =const. Let us choose the Riemannian fibre metric on $Y \to \mathbb{R}$ which takes the form

$$g_{y'y'} = \exp(2\lambda t)$$

with respect to the coordinate y'. Then relative to the coordinate y, it reads

$$g_{yy} = \frac{\partial y'}{\partial y}\frac{\partial y'}{\partial y} g_{y'y'} = \exp(2(\lambda - 1)t). \tag{6}$$

The instantwise distance between the solutions s and s' (5) with respect to the metric g (6) is

$$\rho_t(s, s') = [g_{yy}(s(t) - s'(t))^2]^{1/2} = |c - c'| \exp(\lambda t).$$

One at once obtains that the Lyapunov spectrum of any solution of the differential equation (3) with respect to the metric (6) is λ.